

TEV EMISSION FROM THE GALACTIC CENTER BLACK-HOLE PLERION

ARMEN ATOYAN¹ AND CHARLES D. DERMER²¹ CRM, Université de Montréal, C.P. 6128, Montréal, Canada H3C 3J7; atoyan@crm.umontreal.ca² E.O. Hulburt Center for Space Research, Code 7653, Naval Research Laboratory, Washington, DC 20375-5352; dermer@gamma.nrl.navy.mil

Draft version February 2, 2008

ABSTRACT

The HESS collaboration recently reported highly significant detection of TeV γ -rays coincident with Sgr A* (Aharonian et al. 2004). In the context of other Galactic Center (GC) observations, this points to the following scenario: In the extreme advection-dominated accretion flow (ADAF) regime of the GC black hole (BH), synchrotron radio/sub-mm emission of ~ 100 MeV electrons emanates from an inefficiently radiating turbulent magnetized corona within $20 R_S$ (Schwarzschild radii) of the GCBH. These electrons are accelerated through second-order Fermi processes by MHD turbulence, as suggested by Liu et al. (2004). Closer to the innermost stable orbit of the ADAF, instabilities and shocks within the flow inject power-law electrons through first-order Fermi acceleration to make synchrotron X-ray flares observed with Chandra, XMM, and INTEGRAL. A subrelativistic MHD wind subtending a ~ 1 sr cone with power $\gtrsim 10^{37}$ erg s⁻¹ is driven by the ADAF from the vicinity of the GCBH. As in pulsar powered plerions, electrons are accelerated at the wind termination shock, at $\gtrsim 10^{16.5}$ cm from the GCBH, and Compton-scatter the ADAF and the far infra-red (FIR) dust radiation to TeV energies. The synchrotron radiation of these electrons forms the quiescent X-ray source resolved by Chandra. The radio counterpart of this TeV/X-ray plerion, formed when the injected electrons cool on timescales of $\gtrsim 10^4$ yrs, could explain the origin of nonthermal radio emission in the pc-scale bar of the radio nebula Sgr A West.

Subject headings: acceleration of particles—black hole physics—Galaxy: center—plasmas—radiation processes: nonthermal

1. INTRODUCTION AND OBSERVATIONS

HESS measurements of TeV fluxes from Sgr A* (Aharonian et al. 2004) confirm, with high significance, earlier reports by the Whipple (Kosack et al. 2004) and CAN-GAROO (Tsuchiya et al. 2004) collaborations (albeit at higher flux levels than HESS) about TeV emission from the Galactic Center. This result adds important new information about the massive ($M_{GC} \approx 4 \times 10^6 M_\odot$; Ghez et al. (2003)) BH at the center of the Galaxy. When combined with the wealth of observational data in the radio to X-ray wavebands (for review, see Melia & Falcke 2001), the TeV γ -ray data point to a scenario involving an inner, inefficiently radiating magnetized corona, i.e. the ADAF, and a subrelativistic outflow of particles and field from the ADAF. As in pulsar plerions like Crab Nebula (see Kennel & Coroniti (1984)), the wind terminates at a shock, at $R \gtrsim 3 \times 10^{16}$ cm from the GCBH, to create a black-hole powered ‘plerion’ where the TeV radiation is produced.

The VLBI size of the radio source at the location of Sgr A* is $110(\pm 60) \mu$ as at $\nu = 215$ GHz and $190(\pm 30) \mu$ as at $\nu = 87$ GHz (Krichbaum et al. 1998). Using VLBA, Bower et al. (2004) resolve Sgr A* at 42 GHz to have an angular size of $240(\pm 10) \mu$, or $R_{rad} \simeq 24 R_S$ for $M_{GC} = 4 \times 10^6 M_\odot$ black hole and a GC distance $d = 8$ kpc, and find that the size increases with wavelength. The radio emission of Sgr A* is rather stable, with total luminosity $L_{36} = L_{rad}/10^{36}$ erg s⁻¹ $\simeq 1$ and spectral flux $S_\nu \propto \nu^{0.3}$ that sharply falls off above $\nu_0 \sim 10^{12}$ Hz (Zylka et al. 1995; Falcke et al. 1998).

The near-infrared (NIR) flares show factors-of-2 flux variations on timescales of ~ 20 min and reach powers of $\approx 10^{35}$ erg s⁻¹ (Genzel et al. 2003). The spectra rise in νS_ν diagrams, and show powers $\approx 10^{35}$ erg s⁻¹, similar with the (*nonsimultaneously*) detected X-ray flares (Baganoff et al. 2001; Porquet et al. 2003). The 200 s X-ray variability timescale limits the engine size to $r = R/R_S \lesssim 5$. The unabsorbed quiescent flux detected by Chandra observatory from Sgr A* is $L_{X,qui}(2-10 \text{ keV}) \approx 2.4 \times 10^{33}$ erg s⁻¹, and has angular extent $\theta_{FWHM} \approx 1.4''$ (Baganoff et al. 2003), corresponding to source size $R_X \approx 8.3 \times 10^{16}$ cm.

The TeV spectral fluxes in two HESS campaigns in June/July and July/August of 2003 are almost exactly the same, extending with photon index $\approx 2.1-2.3$ from 160 GeV to 10 TeV, with total power $L_\gamma \approx 10^{35}$ erg s⁻¹. There is no sign of flux variation within the statistics available in each of these sets (Aharonian et al. 2004). Given the high statistical significance of the signal in both data sets, these results suggest a quiescent or quasi-stationary TeV source coincident within $\leq 1.5'$ of Sgr A*.

We propose a model for Sgr A* where the quasi-stationary radio and flaring X-ray/NIR emissions are synchrotron radiation from the ADAF. The MHD wind from the ADAF powers a black-hole plerion where the quiescent X-ray and TeV emissions are produced by electrons accelerated at the wind shock. The model explains the existing multi-wavelength data, and makes predictions that can be checked by various X-ray and gamma-ray detectors.

2. PARAMETERS OF THE ADAF

The dimensionless mass-accretion rate (or Eddington ratio) $\dot{m} = \eta_{BH} \dot{M} c^2 / L_{Edd}$ is defined in units of the Eddington luminosity L_{Edd} , where $\eta_{BH} \simeq 0.1$ is the maximum efficiency of gravitational-to-photon energy conversion by the BH. In the ADAF model (Narayan & Yi 1995; Esin et al. 1998), thermal radiation of the accretion plasma at the level $L_T = \dot{m}^2 L_{Edd} / \dot{m}_*$ is predicted when $\dot{m} \leq \dot{m}_* \sim 0.1$. The rate $\dot{m} \simeq 1.5 \times 10^{-5}$ is obtained if one formally equates L_T with the observed $L_{rad} \simeq 10^{36} \text{ erg s}^{-1}$.

The compact radio emission cannot originate from an optically-thick accretion disk which would only be allowed at $R \gg R_{rad}$, or from a much hotter ADAF plasma at smaller scales, but is compatible with a synchrotron origin (Liu, Petrosian, & Melia 2004). The thermal output of the ADAF would be peaked mostly in the hard X-ray/soft gamma-ray domain, where the quiescent luminosity is at least 2 orders of magnitude smaller than L_{rad} . This results in the accretion rate estimate $\dot{m}_{-6} = \dot{m} / 10^{-6} \lesssim 10^{-6}$ consistent with ADAF model and Sgr A* observations. Even for $\dot{m}$ this small, $\lesssim 10^{39} \text{ erg s}^{-1}$ of accretion power is still either advected into the BH or escapes as an outflow.

Adopting GCBH mass $3 \times 10^6 M_\odot$, the ADAF magnetic field at radii $r = R/R_S$ is (Narayan & Yi 1995)

$$B(r) \approx 370 \alpha^{-1/2} \lambda^{1/2} \dot{m}_{-6}^{1/2} r^{-5/4} \text{ G}, \quad (1)$$

where $\lambda = 1 - \beta$ is the ratio of the magnetic to the total pressure in the accreting gas and α is the viscosity parameter. Using $\alpha \sim 0.1$ and $\lambda \sim 0.15$ (Quataert & Narayan 1999), the magnetic field $B \sim 10 \text{ G}$ in the ADAF at $r \lesssim 20$ is found. For these fields, the Lorentz factor of electrons producing synchrotron radiation peaked at $\nu_0 \gtrsim 10^{12} \text{ Hz}$ is $\gamma_0 \sim 200$, and the ADAF is still transparent to self-absorption for $\gtrsim \text{GHz}$ radiation.

A stochastic acceleration model for the radio-mm emission from Sgr A* (Liu, Petrosian, & Melia 2004) can be quantified using the second-order Fermi acceleration theory of Dermer, Miller, & Li (1996). The magnetic field is defined through the relation $B^2/8\pi = \epsilon_B \eta_{BH} \dot{M} c^2 / (4\pi R^2 c)$, implying $B \approx 30 \epsilon_B^{1/2} L_{36}^{1/4} \text{ G}$ for a region of size $20 R_S$. Equating the acceleration rate of electrons by whistler turbulence with the synchrotron loss rate gives a characteristic electron Lorentz factor $\gamma_0 \cong 5 (\zeta \epsilon_B)^{1/3} L_{36}^{1/6} \tau_T^{-11/18}$, assuming a Kolmogorov spectrum for the turbulence. Here ζ is the fractional whistler turbulent energy density compared to $B^2/8\pi$, and τ_T is the Thomson depth of the ADAF, assumed to be composed of e-p plasma. Values of $B = 10 \text{ G}$ and $\gamma_0 \cong 200$ are implied for the ADAF when $\zeta \approx 0.1$, $\epsilon_B \approx 0.1$ and $\tau_T \approx 2 \times 10^{-4}$.

3. BLACK HOLE PLERION

A Compton (i.e., leptonic) origin for non-variable TeV flux detected with HESS implies that the synchrotron counterpart of that flux is the quiescent component of the X-ray flux. The magnetic-field energy density in the source should then be $u_B = B^2/8\pi < 0.1 u_{rad}$, where

$$u_{rad} = \frac{L_{rad}}{4\pi c R^2} = 2.65 \times 10^{-8} \frac{L_{36}}{R_{16}} \text{ erg cm}^{-3} \quad (2)$$

is the energy density of the target photons at $R = R_{16} 10^{16} \text{ cm}$. Inside the radio source, at $r \lesssim 20$, the radiation energy density saturates at the level $u_{rad} \simeq L_{rad}/2\pi c R_{rad}^2$. Thus, the origin of TeV flux at those small

distances would require magnetic fields $B \sim 0.1 \text{ G}$, far below the ADAF model values.

In the ADIOS (“advection-dominated inflow-outflow solution” Blandford & Begelman (1999)) extension of the ADAF model, the outflow may carry a significant fraction of the generated kinetic energy of the plasma from the BH vicinity to large distances. We assume a wind of magnetized accretion plasmas with total power $L_{wind} = L_{w.37} 10^{37} \text{ erg s}^{-1}$ propagating in a two-sided cone with opening angle Ω at speed $v_w \lesssim c$; for calculations we use $v_w = c/2$. This wind terminates at a subrelativistic shock at $R_{shock} \leq R_X$ from the GCBH. The shock accelerates electrons to energies $E_{max} = \gamma_{max} m_e c^2 \gg 10 \text{ TeV}$ to produce the observed TeV γ -rays by Compton upscattering of sub-mm ($\nu_0 \sim 10^{12} \text{ Hz}$) photons from the ADAF. Another important photon target is the emission from the cold ($\simeq 100 \text{ K}$) dust ring of Sgr A West, with total luminosity $\simeq 5 \times 10^6 L_\odot$ in the central arcmin (few pc) region around Sgr A* (Becklin et al. 1982). This radiation has a spectral peak at $\simeq 10^{13} \text{ Hz}$ and estimated energy density $\simeq 2.4 \times 10^3 \text{ eV cm}^{-3}$. It becomes the main contributor to the $\lesssim 100 \text{ GeV}$ Compton flux from the plerion at $R \gtrsim 10^{17} \text{ cm}$ from Sgr A*. Synchrotron radiation from these same multi-TeV electrons at $R \leq 10^{17} \text{ cm}$ produces the quiescent X-ray flux from Sgr A*.

Chandra observations (Baganoff et al. 2003) reveal diffuse thermal X-rays with luminosity $2.4 \times 10^{34} \text{ erg s}^{-1}$ from the central parsec region of Sgr A*, produced by $kT \simeq 1.3 \text{ keV}$ plasma with $n_e = 26/\eta_f^{1/2} \text{ cm}^{-3}$ density, where $\eta_f = \eta_{-1}/10^{-1}$ is the volume filling factor. Equating the gas pressure $n_e kT$ in that region with the energy density of the wind gives the shock distance

$$R_{shock} \simeq 3.1 \times 10^{16} L_{w.37}^{1/2} \Omega_w^{-1/2} \eta_{-1}^{1/4} \text{ cm}. \quad (3)$$

The TeV luminosity $L_\gamma \simeq 10^{35} \text{ erg s}^{-1}$ implies a total electron acceleration power $L_e \gtrsim 10^{36} \text{ erg s}^{-1}$. Assuming that the efficiency of electron acceleration $L_e/L_{wind} \lesssim 30\%$, and letting the rest go to thermal plasma and nonthermal protons (unlike in pulsar winds where relativistic electrons dominate), we arrive at $L_{w.37} \gtrsim 0.3$. Note that Eq.(3) predicts that the maximum power of the outflow sustainable at distances $R_{shock} \leq R_X$ is $L_{w.37} \lesssim 10$.

The mean fluid speed v_f in the TeV plerion formed downstream of the shock can be derived from $(R_X - R_{shock}) \simeq v_f \times t_C(\gamma_X)$, where

$$t_C(\gamma_X) \simeq 4 \times 10^7 \frac{R_{16}^2}{L_{36}} \left(\frac{\epsilon_X}{1 \text{ keV}} \right)^{-1/2} \text{ s} \quad (4)$$

is the Compton cooling time of electrons producing synchrotron X-rays with energy ϵ_X for the estimated magnetic field $B_{pler} \sim 100 \mu\text{G}$ in the GCBH plerion at scales $R \lesssim R_X$. Taking $R_{16} = 5$ for the median distance of the TeV plerion, we find $v_f \gtrsim 500 \text{ km/s}$ at those distances.

Since relativistic electrons of all energies convect with the fluid at the same speed, the TeV plerion model predicts shrinking of the quiescent X-ray source size at higher energies. Note also that the opening angle of the outflow should be $\Omega \sim 1$ rather than $\Omega \ll 1$, since otherwise R_{shock} would exceed R_X . This indicates that the energy outflow is in the form of a wind rather than well-collimated jet.

4. SPECTRAL MODELING

Fig. 1 shows numerical calculations of the quiescent radiation components of Sgr A*. These include synchrotron and SSC emission from the magnetized ADAF within $r_{rad} \approx 20$ of the GCBH (*dashed curves*), X-ray and TeV emission from the compact plerion at scales $R \sim (3 - 10) \times 10^{16}$ cm (*solid curves*), and the emission from the larger plerion (*dot-dashed curves*) inflated to *pc* scales in the process of convective (and possibly also diffusive) propagation of the accelerated electrons on timescales $\lesssim 10^4$ yr. Magnetic field in the latter, $B_2 = 140 \mu\text{G}$, is assumed higher than $B_1 = 90 \mu\text{G}$ in the TeV plerion. This is in agreement with the expected increase of the magnetic field downstream of the shock in typical pulsar plerions (Kennel & Coroniti 1984), and is explained by deceleration and compression of the plasma. But it is also possible that $B_2 < B_1$, in which case a smaller synchrotron flux from the *pc*-scale plerion than shown in Fig. 1 is predicted.

The inset in Fig. 1 shows the nonthermal electron distributions. Besides synchrotron and Compton losses, calculations also include Coulomb and bremsstrahlung losses in a medium with $n_{gas} = 10^3 \text{ cm}^{-3}$ characteristic for central *pc* regions of Sgr A*. Electrons with a power-law injection index $\alpha_{pler} = 2.2$ and an exponential cutoff above $E_{max} = 50 \text{ TeV}$ are injected into 2-sided nebula with total power $L_e = 6.5 \times 10^{36} \text{ erg s}^{-1}$. These electrons radiatively cool downstream of the shock and propagate to distances $\geq 10^{17}$ cm on the timescale $t_{esc} = 50 \text{ yr}$, during which most of the injected energy of the multi-TeV electrons (producing TeV and X-ray fluxes observed) has been already lost.

The radio emission is produced at $r \lesssim 20$ in $B \approx 10 \text{ G}$ field by a relativistic Maxwellian electron distribution $\propto \gamma^2 e^{-\gamma/\gamma_0}$ with $\gamma_0 = 200$ resulting from a balance between second-order energy gains and synchrotron losses (Schlickeiser 1985). The stochastic acceleration power of radio electrons is therefore about equal to the observed radio luminosity L_{rad} .

The SSC emission component of these electrons, shown in Fig. 1 by the dashed curve, falls in the sub-keV region below the level of the quiescent X-ray flux. However, if magnetic fields $\lesssim 5 \text{ G}$ are assumed, this SSC contribution would rise and also move to 1 keV, and could contribute to radiation at that energies. It cannot, however, explain the entire quiescent flux extending to 10 keV.

The powerful X-ray and NIR flares observed from Sgr A* are explained in our model by the onset of instabilities of the accretion flow at distances of only few R_S . They result in strong shocks in the accretion flow and effective acceleration of particles advected with the flow. The accelerated particles are then easily taken out of this region in the wind/jet outflow. Synchrotron emission of electrons accelerated to $\gamma > 10^6$ explains the X-ray flares with very short variability scales. Furthermore, while propagating through the first few tens of R_S , these electrons have sufficient time, $\sim 10^3 \text{ s}$, to cool in the high B fields there down to $\gamma \lesssim 3 \times 10^3$ to produce flares in the NIR domain. Self-absorbed flares at $\lesssim 100 \text{ GHz}$ detected on $\lesssim 1 \text{ day}$ timescales after the X-ray flares (Zhao et al. 2004) could be explained by the radiation from these same electrons at later stages/larger distances of the outburst in the ‘expanding source’ scenario.

In Fig. 2 we show the flare fluxes expected in this model.

After shock-acceleration close to the BH and injection into collimated wind outflow with speed $c/2$, electrons propagate through the inner $r \simeq 10\text{--}30$ region on timescale $t_{esc} = 1200 \text{ s}$, after which the radiation losses drop because of wind expansion and decline of B . The solid, dashed, and dot-dashed curves show synchrotron and Compton fluxes produced at times 200 s, 1200 s and 1 hr after the onset of the flare. The electron injection time profile is $L_{e,flare}(t) = L_0/(1 + t/t_0)^2$, with $t_0 = 720 \text{ s}$. The mean magnetic field in ADAF outflow could be enhanced at the flaring state, so we take $B = 25 \text{ G}$. The Compton fluxes shown (thin curves) clearly demonstrate that no detectable TeV flares should be expected during powerful X-ray flares. This is in agreement with the non-detection of TeV flux variations during many days of observation with the HESS telescopes, whereas the X-ray flares occur with frequency of ~ 1 per day (Baganoff et al. 2003).

5. SUMMARY AND CONCLUSIONS

A model consisting of magnetized coronal ADAF (synchrotron) radio emission within $\approx 20 R_S$ of the GCBH, and of a black-hole plerion powered by a wind from the ADAF resolves many puzzling observations of Sgr A*. X-ray flares are synchrotron radiation of electrons accelerated through first-order Fermi process by shocks within a few R_S of the GCBH. Electrons accelerated to $\gamma \gtrsim 10^8$ at the wind termination shock at $\gtrsim 3 \times 10^{16} \text{ cm}$ creates the GCBH plerion. Its synchrotron X-ray emission has been resolved as an $\simeq 1.4''$ source with Chandra (Baganoff et al. 2003). The multi-TeV electrons Compton scatter the radio photons from Sgr A* and FIR photons from the dust ring of Sgr A West to produce the TeV emission. TeV emission from the GCBH plerion is nearly stationary because the cooling time of TeV electrons is $\sim 100 \text{ yrs}$.

A “jet-ADAF” model for Sgr A* has been proposed by Falcke & Markoff (2000) and Yuan, Markoff, & Falcke (2002). Our model, though also based on energy outflow from the ADAF, differs greatly. In particular, the quiescent and flaring X-ray components are produced at different sites. The origin of the X-ray and NIR flares are explained as synchrotron emission of electrons accelerated during episodes of instabilities very close to the BH, not as Compton radiation in the SSC scenario of the “jet-ADAF” model. Importantly, the TeV flux cannot be easily explained if the observed X-rays were due to the SSC mechanism in the vicinity of the GCBH, as suggested by Falcke & Markoff (2000).

Propagation of GeV electrons from the plerion on timescales of $\gtrsim 10^4 \text{ yr}$ with speeds $\sim 100 \text{ km s}^{-1}$ could significantly contribute to the radio synchrotron flux of Sgr A West. In particular, it could explain the suggested nonthermal origin of radiation in the bar of Sgr A West (Wright et al. 1987), formed in the outflow direction.

We also predict that quasi-stationary Compton and bremsstrahlung fluxes from the *pc*-scale plerion, coincident with the central parts of Sgr A West, will be significantly detected and possibly resolved with GLAST at GeV energies. But the expected highly variable Compton counterpart of the synchrotron X-ray flares from the GCBH vicinity is too weak to be detectable with GLAST or HESS. The synchrotron extension of the flare emission in the hard X-ray/soft γ -ray domain explains the $\simeq 40 \text{ min}$

episode of a profound increase of flux detected by INTEGRAL from the direction of Sgr A* (Bélanger et al. 2004).

The absence of apparent TeV flux variations may suggest a proton origin for the TeV radiation (Aharonian & Neronov 2004). Indeed, proton and ion acceleration through first- and second-processes in the ADAF and through first-order acceleration at the wind termination shock could make cosmic rays to produce TeV emission through nuclear pp -interactions with $n \sim 10^3 \text{ cm}^{-3}$ dense gas on pc scales, and could form extended TeV emission possibly already detected with HESS (Aharonian et al. 2004). The hadronic origin of the TeV radiation in the ADAF itself in the BH vicinity requires, however, an extremely dense gas target or extremely large proton powers $\gtrsim 10^{39} \text{ erg s}^{-1}$.

Finally, we note that the unidentified EGRET source 3EG J1746-2851 towards the GC (Mayer-Hasselwander

et al. 1998) is significantly displaced (Dingus & Hooper 2002) from the direction of the GCBH, and is unlikely to be related to Sgr A*. It is probably emission from a young pulsar, though not with the “mouse” PSR J1747-2958 (McLaughlin & Cordes 2003). A young pulsar with γ -ray properties like Vela but with apparent γ -ray power $\approx 10\times$ larger could have been missed in pulsar surveys due to the large dispersion measure towards the GC. We suggest a deeper, higher radio-frequency and X-ray search at the refined location of 3EG J1746-2851.

We thank I. Grenier, S. Markoff, R. Narayan, P. Ray, and R. Romani for discussions. Research of CD and visits of AA to the NRL High Energy Space Environment Branch are supported by *GLAST* Science Investigation No. DPR-S-1563-Y. The work of CD is supported by the Office of Naval Research.

REFERENCES

- Aharonian, F., et al. 2004, *A&A*, 425, L13
 Aharonian, F., and Neronov, A. 2004, *ApJ*, submitted (astro-ph/0408303)
 Baganoff, F. K., et al. 2001, *Nature*, 413, 45
 Baganoff, F. K., et al. 2003, *ApJ*, 591, 891
 Becklin, E. E., Gatley, I., and Werner, M. W. 1982, *ApJ* 258, 135
 Bélanger, G., et al. 2004, *ApJ*, 601, L163
 Blandford, R. D., and Begelman, M. C. 1999, *MNRAS* 303, L1
 Bower, G. C., et al. 2004, *Science*, 304, 704
 Dermer, C. D., Miller, J. A., and Li, H. 1996, *ApJ*, 456, 106
 Dingus, B. & Hooper, D. 2002 COSPAR Plenary Meeting (astro-ph/0212509)
 Esin, A. A., Narayan, R., Cui, W., Grove, J. E., & Zhang, S. 1998, *ApJ*, 505, 854
 Falcke, H., Goss, W. M., Matsuo, H., Teuben, P., Zhao, J.-H., Zylka, R. 1998, *ApJ* 499, 731
 Falcke, H., and Markoff, S. 2000, *A&A* 362, 113
 Genzel, R., Schödel, R., Ott, T., Eckart, A., Alexander, T., Lacombe, F., Rouan, D., & Aschenbach, B. 2003, *Nature*, 425, 934
 Ghez, A. M., et al. 2003, *ApJ*, 586, L127
 Kennel, C. F., & Coroniti, F. V., 1984, *ApJ*, 283, 964
 Kosack, K., et al. 2004, *ApJ*, 608, L97
 Krichbaum, T. P., et al. 1998, *A&A*, 335, L106
 Liu, S., Petrosian, V., & Melia, F. 2004, *ApJ*, 611, L101
 Mayer-Hasselwander, H. A., et al. 1998, *A&A*, 335, 161
 McLaughlin, M. A., & Cordes, J. M. 2003, in *Proc. 4th AGILE Science Workshop* (astro-ph/0310748)
 Melia, F. & Falcke, H. 2001, *ARA&A*, 39, 309
 Narayan, R., and Yi, I. 1995, *ApJ* 452, 710
 Porquet, D., Predehl, P., Aschenbach, B., Grosso, N., Goldwurm, A., Goldoni, P., Warwick, R. S., & Decourchelle, A. 2003, *A&A*, 407, L17
 Quataert, E., and Narayan, R. 1999, *ApJ* 516, 399
 Schlickeiser, R. 1985, *A&A*, 143, 431
 Tsuchiya, K., et al. 2004, *ApJ*, 606, L115
 Yuan, F., Markoff, S., and Falcke, H. 2002, *A&A* 383, 854
 Wright, M. C. H., et al. 1987, in the *Galactic Center* (AIP Conf. Proc. 155), ed. D. C. Backer (New York: AIP), p. 133
 Zhao, J., Herrnstein, R. M., Bower, G. C., Goss, W. M., & Liu, S. M. 2004, *ApJ*, 603, L85
 Zylka, R., Mezger, P. G., Ward-Thompson, D., Duschl, W. J., & Lesch, H. 1995, *A&A*, 297, 83




FIG. 1.— Quasi-stationary/quiescent synchrotron (*heavy curves*) and Compton (*thin curves*) fluxes expected from Sgr A*. Dashed curves show the fluxes from the magnetized ADAF corona with $B = 10 \text{ G}$ and $R \simeq 20 R_S$ due to a relativistic Maxwellian distribution of electrons with mean Lorentz-factor $\gamma_0 = 200$. The solid curves show the fluxes from the BH plerion with $B_1 = 90 \mu\text{G}$ at a distance $R_{\text{pler}} \sim 5 \times 10^{16} \text{ cm}$; the calculations assume escape of the flow from the TeV plerion to $R \geq 10^{17} \text{ cm}$ on timescale 50 yr. The *dot-dashed curves* show the fluxes formed during assumed $t_{\text{esc}} = 2 \times 10^4 \text{ yr}$ propagation time in the plerion at scales $R \sim 0.03 - 3 \text{ pc}$. The *dotted curve* shows the bremsstrahlung from the central few pcs of Sgr A West with gas density $n_{\text{gas}} = 10^3 \text{ cm}^{-3}$. The inset shows the electron energy distributions formed in those 3 spatial scales. The hatched regions in the X-ray domain show the fluxes detected by Chandra ($\alpha_X \sim 2.2 - 3$ are plotted) and XMM ($\alpha_X = 2.5 \pm 0.3$), and the TeV fluxes detected by HESS (see text for references). The radio data are from Zylka et al. (1995) and Falcke et al. (1998).




FIG. 2.— Synchrotron and Compton fluxes (*thick* and *thin* curves, respectively) expected during a flare at 200 s (*solid*), 1200 s (*dashed*), and 3600 s (*dot-dashed* curves) after the onset of the injection of relativistic electrons with $\alpha_{pler} = 2.2$ and $\gamma_{max} = 10^7$, initial injection power $L_e(0) = 6.3 \times 10^{36} \text{ erg s}^{-1}$, and characteristic decline time $t_0 = 720 \text{ s}$ (see text). Escape of electrons with the conical wind during $t_{esc} = 1200 \text{ s}$ from the central $R \leq 20 R_S = 1.8 \times 10^{13} \text{ cm}$ region of the ADAF is supposed. The dotted curve shows the expected bremsstrahlung γ rays at $t = 200 \text{ s}$, calculated for plasma density of 10^8 cm^{-3} in the ADAF.